

Micronucleus Assay of Buccal Mucosa Cells in Waterpipe (Hookah) Smokers: A Cytologic Study

Mehdi DehghanNezhad¹, Noushin Jalayer Naderi², Hassan Semyari³

1. Faculty of Dentistry, Shahed University, Tehran, Iran
2. Department of Oral and Maxillofacial Pathology, Faculty of Dentistry, Shahed University
3. Department of Periodontics, Faculty of Dentistry, Shahed University

KEYWORDS

Assay, Buccal mucosa,
Genotoxicity test,
Micronucleus,
Smoking

Scan to discover online



Main Subjects:
Oral Pathology

Received 14 Jan 2019;

Accepted 23 Feb 2020;

Published Online 26 Feb 2020;

doi [10.30699/ijp.2020.101701.2010](https://doi.org/10.30699/ijp.2020.101701.2010)

ABSTRACT

Background & Objective: Micronucleus assay of buccal mucosa cells is a simple bio-monitoring method for diagnosing the genetic damages of toxic agents. The aim was to study the genotoxic effect of waterpipe smoking on buccal mucosa cells using micronucleus assay.

Methods: This was a case control. A total of 30 male waterpipe smokers and 30 non-smokers were included in the study. The exfoliated buccal mucosa cells were scrapped using wooden spatula and were spread over glass slides. The mean number of micronuclei was determined using Feulgen-stained slides. The number of micronuclei per 1000 cells was calculated and compared between the two groups of smokers and non-smokers.

Results: The mean number of micronuclei in waterpipe smokers and non-smokers was 1.94 ± 0.39 and 1.68 ± 0.35 , respectively. The micronuclei count in waterpipe smokers was significantly higher than non-smokers ($P=0$). The difference between the number of waterpipe smoking and micronuclei count was significantly different ($P=0$).

Conclusion: The mean number of micronuclei in buccal mucosa cells of waterpipe smokers was significantly higher than non-smokers. The genotoxicity effect of waterpipe was dose-dependent.

Corresponding Information:

Noushin Jalayer Naderi, Department of Oral and Maxillofacial Pathology, Faculty of Dentistry, Shahed University, Tehran, Iran Email: jalayer@shahed.ac.ir

Copyright © 2020. This is an open-access article distributed under the terms of the Creative Commons Attribution- 4.0 International License which permits Share, copy and redistribution of the material in any medium or format or adapt, remix, transform, and build upon the material for any purpose, even commercially.

Introduction

For many centuries, waterpipe is used for tobacco smoking in Asia and Africa. Traditionally, most waterpipe users are concentrated in North Africa and South-east Asia (1). Nowadays, waterpipe smoking is a global problem. Based on published data numbers of waterpipe users are increasing among women and teenagers. Waterpipe smoking delivers high levels of nicotine to mucosal cells of oral cavity and respiratory tract. The inhaled nicotine contains toxic materials comprising carbon monoxide and carcinogenic materials (2). The risk of carcinomatous changes intensifies by waterpipe smoke (3-4).

Waterpipe smokers inhale higher doses of nicotine compared to a cigarette smoker. Inhalation a chemical agents such as nicotine causes genetic damages. Revealing the genetic damage in persons who are at risk on being exposed to toxic materials is a practical tool in evaluating the genotoxicity effect of agents and malignant transformation. Bio-monitoring of individuals exposed to genotoxic agents using exfoliated buccal mucosa cells is a simple and a reliable

method for determining the genotoxic effect. For the first time, Stich et al. used the micronucleus test on exfoliated buccal mucosa cells for tracing the genotoxic exposure in humans (5). Micronucleus test is an inexpensive and non-invasive method for screening the persons who are at risk of cancer development (6). Micronucleus is a separated part of nucleus originates during cellular division. Micronuclei generate from chromosomal fragments of inter-phasic cells (7). The micronuclei are cytoplasmic structures measuring between 1/5 to 1/3 size of nucleus with staining similar to nucleus (8). In general populations, the mean prevalence of cells with micronuclei is 0 to 0.9%. Any increase in micronucleus count is a reflection of chromosomal alterations. The number of micronuclei has been related to degree of carcinogenic effect (9).

In a study which was among the first investigations about the effect of waterpipe smoking on cytogenetic changes, El-Setouhy et al. showed a higher level of micronuclei in waterpipe smokers of rural Egypt population (10). In Iran, the popularity of waterpipe

smoking is growing. This is an important issue for persons who are concerned about health planning programs. Despite increasing tendency of youths and women for using the waterpipe, knowledge about the genotoxicity of waterpipe is insufficient. The aim was to evaluate the genotoxic effects of waterpipe smoking by testing the micronucleus count of buccal mucosa cells in a cytologic study.

Materials and Methods

This was a case control with simple sampling method. The study was carried out in the department of Pathology, Faculty of Dentistry, Shahed University, Tehran, Iran in Oct 2015-Apr 2016. The study was approved by ethical committee of Shahed University and registered as IR.Shahed.Rec.1394.301.

Using Cochran's sample size formula with 95% confidence level and 90% strength of test, the sample size was determined as 27.89 subjects in both smokers and non-smokers' groups. A total 60 subjects (30 waterpipe smokers and 30 non-waterpipe and cigarette smokers) were entered the study. All subjects in both case and control groups were 20 to 50 year old males. The persons younger than 20 years old, suffering of systemic disease and any oral lesions, consuming any type of drugs and being exposed to dental radiography beam in recent 6 months and alcohol consumers were excluded from the study in both case and control groups. Waterpipe smokers were selected from a local waterpipe café in Tehran, Iran. Non-smokers were collected from dental school of Shahed University. All subjects were living in Tehran and were not farmer or worker in Arsenic industries.

The inclusion criteria for selecting the waterpipe users were using the waterpipe at least once in a week. To reduce the effect of cigarette smoking on the results, the protocol of El-Setouhy *et al.* was used to select the samples. Accordingly, persons who never smoked cigarettes or smoke utmost 100 cigarettes in their whole life were included the study (10). Time duration of waterpipe smoking was registered based on number of smokings per year (11).

An inform consent was taken from all subjects before participation in the study. The demographic information were entered in a registration form and coded. The participants were not identified by names and families.

For collecting the buccal mucosa cells, all subjects rinsed their mouth twice with normal saline. Using wooden spatula, the exfoliated buccal cells were scrapped and were spread on to the glass slides. Samples were fixed in Carnoy's fixative (methanol and glacial acetic acid in a ratio of 3:1) for 30-35 minutes and then dried at room temperature. The modified method of Thomas *et al.* was used for staining the micronuclei by Feulgen reaction (12). The Feulgen reaction was performed as follows: Slides were dipped in 1 N HCL at 60°C for 10 minutes, rinsed in the distilled water for 3 minutes, placed in Schiff's reagent

for 90 minutes and then in normal saline for 10 minutes. Then, slides were placed in 0.5 % sodium metabisulfite solution for 3 times and then rinsed with tap water. Then the slides were stained with 1% light green for 15 minutes, were rinsed with tap water and finally dried and mounted.

The structures within cytoplasm with similar staining of nucleus measuring between 1/5 to 1/3 size of nucleus was considered as micronucleus (8). The cells presenting cell death features comprising of karyorrhexis, karyolysis and pyknosis were not included in the study (Figure 1).

The micronuclei count completed in form of blind. Cells with distinct cellular margin were encountered for counting. The overlapped cells and cellular collections were not considered. Optic microscope (ZEISS, Germany) under oil immersion lens with $\times 1000$ magnifications was used for micronuclei count. The micronuclei count was demarcated by the number of counted micronuclei per 1000 cells per subject (10). Mean number of micronuclei were determined for all samples and were presented as mean \pm SD. The linear regression and T-test were employed at the $P\leq 0.01$ as the significant level. The statistical analyses were completed using SPSS 20 package (IBM Company, Chicago, IL, USA).

Results

The average age of waterpipe smokers and non-smokers were 26.83 ± 3.74 and 28 ± 7.88 years, respectively. The range of waterpipe smoking duration was from 1 to 11 years with the mean duration of 3.3 ± 2.24 years. The mean number of micronuclei in buccal mucosa of waterpipe smokers and non-smokers were 1.94 ± 0.39 and 1.68 ± 0.35 , respectively. The count of micronuclei in buccal mucosa of waterpipe smokers and non-smokers were 25 ± 1.83 and 8.78 ± 0.83 , respectively (Table 1).

The T-test revealed that micronuclei count in waterpipe smokers was significantly higher than non-smokers ($P=0$). The difference between the number of waterpipe smoking and micronuclei count was significantly different ($P=0$) (Figure 2). The comparison of data on number of waterpipe smoking per year using regression analysis indicated that, the number of micronuclei count increased to 0.33 ($P=0.35$) by increasing the smoking time in each years of smoking. Each time waterpipe smoking was associated with an increase in micronucleus count up to 0.027 ($P=0$).

Table 1. The demographic characteristics and micronucleus assay of the waterpipe smokers (n=30) and non-smokers (n=30)

Characteristics	Waterpipe	Smoker	Non	Smoker	Sig (two-tailed)
	Number	Mean± SD	Number	Mean± SD	T-test
Age					
20-30	25	25.5±2.3	23	23.7±2.7	P=0.46
31-40	5	33.2±15.2	5	37.1±3.37	
41-50	0	0	2	44±1.4	
Time duration of smoking*					
0-200	11	118±33.4	0	0	
201-400	15	260±48.1	0	0	
401-600	4	555±52.5	0	0	
MN** count per subject					
0-10	1	8	12	7.5±1.31	P=0
11-20	10	14.8±5.35	17	15.4±2.85	
21-30	10	25.5±3.95	1	21	
31-40	7	38±1.67	0	0	
41-50	2	41	0	0	
Mean per subject MN					
0-1.5	5	1.35±0.08	13	1.37±0.10	P=0
1.6-2	12	1.7±0.15	12	1.75±0.13	
2.1-2.5	10	2.19±0.14	5	2.28±0.20	
2.6-3	3	2.6±0.06	0	0	

* Number of smoking per year

**MN: Micronucleus



Fig. 1. Photomicrograph of a cell with 3 micronuclei (arrows) in buccal mucosa smear of waterpipe smoker ($\times 1000$, Feulgen staining)




Fig. 2. The mean number of micronuclei count in waterpipe smokers and non-smokers

Discussion

The study showed that the mean number of micronuclei in buccal mucosa cells of waterpipe smokers was significantly higher than non-smokers. The genotoxic effect of waterpipe on buccal cells and peripheral blood leukocytes have been demonstrated with comet assay (13), sister chromatid exchanges (SCEs) assay (14) and chromosome analyses (15). In present study, using a simpler method, the previous results were confirmed. Micronucleus assay is a reliable, simple and inexpensive biological test in demonstrating the genotoxic effect of agents. The results of present study showed that the micronuclei count of buccal mucosa cells in waterpipe smokers was higher than non-smokers. The finding is in consistent with El-Setouhy, *et al.* (10).

It has been reported that the micronuclei count in tobacco chewer and cigarette smokers were higher than general population. Based on reports, the count of micronuclei in smokers were 1-2 times more than non-smokers (16-20). Compatible with previous findings in cigarette smokers, the results showed that the mean count of micronuclei in waterpipe smokers is almost 1.5 fold more than persons who never smoked waterpipe.

The false believe about of waterpipe smoking being harmless in comparison to cigarette smoking derives from the method of waterpipe application; passing the smoke of burned tobacco through the water. Smoke of waterpipe contains toxic material such as carbon monoxide and heavy metals (2). The carbon monoxide in expired air of waterpipe and cigarette smoking are 23.7 ppm and 2.7 ppm, respectively. The carboxyhemoglobin level after waterpipe smoking is 3 times higher than cigarette (21).

The amount of produced toxin during one session of waterpipe smoking is equal to 10 cigarettes per day (22). The difference originates from different exposing time to smoke. The average time of cigarette and waterpipe smoking are 5-7 minutes and 45 minutes, respectively. A person inhales 0.5 -0.6 L smoke during cigarette smoking. This amount equals to 0.15 -1 L during waterpipe smoking (1,23).

The present study showed that by adding one year to waterpipe smoking history, the number of micronuclei count increased to 0.33. Each time waterpipe smoking was associated with an increase in micronucleus count up to 0.027. The results were compatible with this finding that the hazard of waterpipe smoking depends on amount and time of smoking (24).

Absence of an established protocol for measuring the dose and duration of waterpipe smoking is a problematic concern in studying the impact of waterpipe smoking on micronucleus assessment. Different staining method has been used in evaluating the micronucleus assessment. Application of the nonspecific DNA stains in demonstrating the micronuclei of epithelial cells leads to false-positive or false-negative results. It has been shown that the results of micronuclei evaluation in oral mucosa cells strongly relates to staining method (25). Omitting the effect of staining method on obtained results, we used the Feulgen stain. Feulgen technique is the most reliable method for staining the nuclear DNA and micronuclei evaluation in cytologic materials (26).

It has been reported that air pollution, exposing to agricultural pesticides and chronic occupational exposure to Arsenic are relating factors in producing the higher rates of micronuclei count in buccal mucosa and peripheral blood lymphocytes (10,27).

In present study, all samples were collected from a local waterpipe café in Tehran. All subjects were under the same condition regarding the inhalation of polluted air. The subjects were not farmer or worker in Arsenic industries. To achieve more reliable results and omit the possible effect of female hormones on findings, the study was completed on 20 to 50 year old males. The used sampling method and using specific DNA stain decreased any possible biases in achieved results.

The present study was limited on male waterpipe smokers. Study the genotoxic effects of waterpipe smoking in females and comparing the results with male users is intensely recommended. In most societies such as Iran, waterpipe smoking is a fun activity. Most waterpipe smokers are not cigarette smokers because they believe that cigarette smoking is more harmful than waterpipe smoking. Alternatively, some waterpipe users are heavy cigarette smokers. Because of this divergence, tissue sampling was very time consuming and difficult. To date, waterpipe smoking is a public health problem. Recent studies showed the increasing level of waterpipe smoking between youths and educated persons (28). Waterpipe smoking has adverse health effects similar to or even higher than cigarette smoking (29). Update socio-demographic researches on waterpipe smoking is an important necessity for managing the preventive efforts.

Conclusion

The waterpipe smoking had genotoxic effect on human buccal mucosa cells. The genotoxic effect of waterpipe was dose-dependent. Due to increasing interest of youths and women in waterpipe smoking, further researches were needed for studying the health effect of waterpipe smoking on different human cells and tissues in both genders and different age ranges.

Acknowledgements

The study was completed by financial support of Shahed University. The author thanks Dr. Mohammad Javad KharaziFard for statistical analyzing.

Conflict of Interest

The authors declared no conflict of interest.

References

- Shihadeh A, Azar S, Antonios C, Haddad A. Towards a topographical model of narghile water-pipe café smoking: a pilot study in a high socioeconomic status neighborhood of Beirut, Lebanon. *Pharmacol Biochem Behav.* 2004; 79:75-82. [DOI:10.1016/j.pbb.2004.06.005] [PMID]
- Maziak W, Ward KD, Afifi Soweid RA, Eissenberg T. Tobacco smoking using a waterpipe: a re-emerging strain in a global epidemic. *Tob Control.* 2004; 13:327-33. [DOI:10.1136/tc.2004.008169] [PMID] [PMCID]
- Nafae A, Misra SP, Dhar SN, Shah SN. Bronchogenic carcinoma in Kashmir Valley. *Indian J Chest Dis.* 1973; 15:285-95.
- El-Hakim IE, Uthman MA. Squamous cell carcinoma and keratoacanthoma of the lower lip associated with "Goza" and "Shisha" smoking. *Int J Dermatol.* 1999; 38:108-10. [DOI:10.1046/j.1365-4362.1999.00448.x] [PMID]
- Stich HF, Curtis JR, Parida BB. Application of the micronucleus test to exfoliated cells of high cancer risk groups: tobacco chewers. *Int J Cancer.* 1982; 30:553-9. [DOI:10.1002/ijc.2910300504] [PMID]
- Chatterjee S, Dhar S, Sengupta B, Ghosh A, De M, Roy S, et al. Cytogenetic monitoring in human oral cancers and other oral pathology: the micronucleus test in exfoliated buccal cells. *Toxicol Mech Methods.* 2009; 19: 427-33. [DOI:10.1080/15376510903127530] [PMID]
- Ramirez A, Saldanha PH. Micronucleus investigation of alcoholic patients with oral carcinomas. *Genet Mol Res.* 2002; 1:246-60.
- Sarto F, Finotto S, Giacomelli L, Mazzotti D, Tomanin R, Levis AG. The micronucleus assay in exfoliated cells of the human buccal mucosa. *Mutagenesis.* 1987; 2:11-7. [DOI:10.1093/mutage/2.1.11] [PMID]
- Stich HF, Rosin MP, Vallejera MO. Reduction with vitamin A and beta-carotene administration of proportion of micronucleated buccal mucosal cells in Asian betel nut and tobacco chewers. *Lancet.* 1984; 1:1204-6. [DOI:10.1016/S0140-6736(84)91692-1]
- El-Setouhy M, Loffredo CA, Radwan G, Abdel Rahman R, Mahfouz E, Israel E, et al. Genotoxic effects of waterpipe smoking on the buccal mucosa cells. *Mutat Res.* 2008; 655:36-40. [DOI:10.1016/j.mrgentox.2008.06.014] [PMID] [PMCID]
- Klokkevold PR, Newman MG, Takei HH, editors. *Carranza's clinical periodontology.* Elsevier Saunders; 2015. P.179.
- Thomas P, Hecker J, Faunt J, Fenech M. Buccal micronucleus cytome biomarkers may be associated with Alzheimer's disease. *Mutagenesis.* 2007; 22:371-9. [DOI:10.1093/mutage/gem029] [PMID]
- Al-Amrah HJ, Aboznada OA, Alam MZ, ElAssouli MZ, Mujallid MI, ElAssouli SM. Genotoxicity of waterpipe smoke in buccal cells and peripheral blood leukocytes as determined by comet assay. *Inhal Toxicol.* 2014; 26: 891-6. [DOI:10.3109/08958378.2014.970787] [PMID]
- Khabour OF, Alsatari ES, Azab M, Alzoubi KH, Sadiq MF. Assessment of genotoxicity of waterpipe and cigarette smoking in lymphocytes using the sister-chromatid exchange assay: a comparative study. *Environ Mol Mutagen.* 2011; 52:224-8. [DOI:10.1002/em.20601] [PMID]
- Derici Eker E, Koyuncu H, Şahin NÖ, Yüksel A, Berköz M, BudakDiler S, et al. Determination of genotoxic effects of hookah smoking by micronucleus and chromosome aberration methods. *Med Sci Monit.* 2016 21; 22:4490-4494. [DOI:10.12659/MSM.898593] [PMID] [PMCID]

16. Stich HF, Stich W, Parida BB. Elevated frequency of micronucleated cells in the buccal mucosa of individuals at high risk for oral cancer: betel quid chewers. *Cancer Lett.* 1982; 17:125-34. [[DOI:10.1016/0304-3835\(82\)90024-6](https://doi.org/10.1016/0304-3835(82)90024-6)] [PMID]
17. Ozkul Y, Donmez H, Erenmemisoglu A, Demirtas H, Imamoglu N. Induction of micronuclei by smokeless tobacco on buccal mucosa cells of habitual users. *Mutagenesis.* 1997; 12:285-7. [[DOI:10.1093/mutage/12.4.285](https://doi.org/10.1093/mutage/12.4.285)] [PMID]
18. Wu PA, Loh CH, Hsieh LL, Liu TY, Chen CJ, Liou SH. Clastogenic effect for cigarette smoking but not areca quid chewing as measured by micronuclei in exfoliated buccal mucosal cells. *Mutat Res.* 2004; 562:27-38. [[DOI:10.1016/j.mrgentox.2004.05.004](https://doi.org/10.1016/j.mrgentox.2004.05.004)] [PMID]
19. Haveric A, Haveric S, Ibrulj S. Micronuclei frequencies in peripheral blood and buccal exfoliated cells of young smokers and non-smokers. *Toxicol Mech Methods.* 2010; 20:260-6. [[DOI:10.3109/15376516.2010.482962](https://doi.org/10.3109/15376516.2010.482962)] [PMID]
20. Konopacka M. Effect of smoking and aging on micronucleus frequencies in human exfoliated buccal cells. *Neoplasma.* 2003; 50:380-2. PMID: 14628093
21. Eissenberg T, Shihadeh A. Waterpipe tobacco and cigarette smoking: direct comparison of toxicant exposure. *Am J Prev Med.* 2009; 37:518-23. [[DOI:10.1016/j.amepre.2009.07.014](https://doi.org/10.1016/j.amepre.2009.07.014)] [PMID] [PMCID]
22. Neergaard J, Singh P, Job J, Montgomery S. Waterpipe smoking and nicotine exposure: a review of the current evidence. *Nicotine Tob Res.* 2007; 9:987-94. [[DOI:10.1080/14622200701591591](https://doi.org/10.1080/14622200701591591)] [PMID] [PMCID]
23. Djordjevic MV, Stellman SD, Zang E. Doses of nicotine and lung carcinogens delivered to cigarette smokers. *J Natl Cancer Inst.* 2000; 92:106-11. [[DOI:10.1093/jnci/92.2.106](https://doi.org/10.1093/jnci/92.2.106)] [PMID]
24. Yadav JS, Thakur S. Genetic risk assessment in hookah smokers. *Cytobios.* 2000; 101(397):101-13.
25. Nersesyan A, Kundi M, Atefie K, Schulte-Hermann R, Knasmüller S. Effect of staining procedures on the results of micronucleus assays with exfoliated oral mucosa cells. *Cancer Epidemiol Biomarkers Prev.* 2006; 15:1835-40. [[DOI:10.1158/1055-9965.EPI-06-0248](https://doi.org/10.1158/1055-9965.EPI-06-0248)] [PMID]
26. Holland N, Bolognesi C, Kirsch-Volders M, Bonassi S, Zeiger E, Knasmueller S, et al. The micronucleus assay in human buccal cells as a tool for biomonitoring DNA damage: the HUMN project perspective on current status and knowledge gaps. *Mutat Res.* 2008; 659:93-108. [[DOI:10.1016/j.mrrev.2008.03.007](https://doi.org/10.1016/j.mrrev.2008.03.007)] [PMID]
27. Vuyyuri SB, Ishaq M, Kuppala D, Grover P, Ahuja YR. Evaluation of micronucleus frequencies and DNA damage in glass workers exposed to arsenic. *Environ Mol Mutagen.* 2006; 47: 562-70. [[DOI:10.1002/em.20229](https://doi.org/10.1002/em.20229)] [PMID]
28. Maziak W, Taleb ZB, Bahelah R, Islam F, Jaber R, Auf R, et al. The global epidemiology of waterpipe smoking. *Tob Control.* 2015; 24 Suppl 1:i3-i12. [[DOI:10.1136/tobaccocontrol-2014-051903](https://doi.org/10.1136/tobaccocontrol-2014-051903)] [PMID] [PMCID]
29. Knishkowsky B, Amitai Y. Water-pipe (narghile) smoking: an emerging health risk behavior. *Pediatrics.* 2005; 116 (1):e113-9. [[DOI:10.1542/peds.2004-2173](https://doi.org/10.1542/peds.2004-2173)] [PMID]

How to Cite This Article

DehghanNezhad, M., Jalayer Naderi, N., Semyari, H. Micronucleus Assay of Buccal Mucosa Cells in Waterpipe (Hookah) Smokers: A Cytologic Study. *Iranian Journal of Pathology*, 2020; (): 75-80. doi: 10.30699/ijp.2020.101701.2010; 15(2):75-80